

# Oleander Aphid, *Aphis nerii* Boyer de Fonscolombe (Insecta: Hemiptera: Aphididae)<sup>1</sup>

Heather J. McAuslane<sup>2</sup>

## Introduction

The oleander aphid, *Aphis nerii* Boyer de Fonscolombe, sometimes called the milkweed aphid, is a common pest of several important ornamental plants in the families Apocynaceae and Asclepiadaceae. This bright yellow aphid with black appendages is commonly found in Florida feeding on oleander, *Nerium oleander*, milkweeds, such as butterfly weed, *Asclepias tuberosa*, and scarlet milkweed, *Asclepias curassavica*, and wax plant, *Hoya carnosa*.

## Distribution

The oleander aphid is cosmopolitan, being found in tropical to warm temperate regions throughout the world. This species probably originated in the Mediterranean region, the origin of its principal host plant, oleander.

## Description

It is thought that the oleander aphid is an obligate parthenogenetic species; thus the adult aphids are all female and males do not occur in the wild. Adult females may be winged or wingless. The winged adult females (alata) are yellow and black with dark wing veins while the wingless forms (apterae) are yellow with black cornicles, antennae, legs, and cauda (tip of abdomen). Nymphs are similar to apterae in appearance except that they are smaller. Size ranges from 1.5 to 2.6 mm in length.



Figure 1. Alata and nymphs of oleander aphid, *Aphis nerii* Boyer de Fonscolombe, on oleander leaf.

Credits: J. Castner, University of Florida

## Life Cycle

Females are viviparous and parthenogenetic, meaning that they deposit nymphs rather than eggs and that the progeny are clones of the adult female (i.e., sexual reproduction is not necessary for offspring production). The nymphs feed gregariously on the plant terminal in a colony that can become quite large. Nymphs progress through five nymphal instars. As in all Sternorrhyncha, there is no pupal stage and adults are produced from the final nymphal instar. Normally apterous adults are produced but alate adults occur under conditions of overcrowding and when plants are senescing, allowing the aphids to migrate to new host

1. This document is EENY247, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date November 2001. Revised January 2012 and May 2017. Visit the EDIS website at <http://edis.ifas.ufl.edu>. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.

2. Heather J. McAuslane, professor; Department of Entomology and Nematology, UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

plants. The parthenogenetic mode of reproduction, high fecundity, and short generation time allow large colonies of oleander aphids to build quickly on infested plants.




Figure 2. Large colony of oleander aphids, *Aphis nerii* Boyer de Fonscolombe, on plant terminal growth.

Credits: Lyle Buss, University of Florida

## Damage

The host range of the oleander aphid includes several genera of Asclepiadaceae (*Gomphocarpus*, *Asclepias*, and *Calotropis*) and Apocynaceae (*Nerium* and *Vinca*). It can occasionally be found infesting plants in the families Compositae, Convolvulaceae, and Euphorbiaceae. In addition, it has been found on citrus. This aphid is able to transmit several viruses including sugarcane mosaic potyvirus and papaya ringspot potyvirus. However, in Florida, the main concern with oleander aphid is the large and unsightly colonies produced on oleander and milkweeds.



Figure 3. Inflorescence of scarlet milkweed heavily infested with oleander aphids, *Aphis nerii* Boyer de Fonscolombe.

Credits: Lyle Buss, University of Florida

The oleander aphid ingests sap from the phloem of its host plant. The damage caused by aphid colonies is mainly aesthetic due to the large amounts of sticky honeydew produced by the colony members and the resulting black sooty mold that grows on the honeydew. In addition, the

growing terminals can be deformed. Of more concern to nursery managers is the potential for stunted plant growth due to repeated heavy infestation throughout the year.



Figure 4. Terminal growth of oleander heavily infested with oleander aphids, *Aphis nerii* Boyer de Fonscolombe.

Credits: Lyle Buss, University of Florida

## Management

Cultural controls offer the best means of managing oleander aphid infestation on oleander. Reduced levels of watering, pruning, and fertilization will reduce the production of tender shoots, the favorite food of oleander aphid. On smaller apocynaceous plants grown as nectar sources for butterflies or as larval host plants for monarch butterflies, the infested shoot can be pruned out and discarded or aphids can be dislodged with a strong stream of water.

Natural biological control can be quite effective in controlling populations of the oleander aphid (Hall and Ehler 1980). The female parasitoid lays eggs in the aphid nymphs. The parasitized aphid develops into a papery, light brown, swollen mummy and the parasitoid develops within this mummy. A single parasitoid emerges from the mummy when the aphid's body has been consumed. In addition, generalist insect predators such as [syrphid larvae](#) in the families Chamaemyiidae, Chrysopidae, and Hemerobiidae and [coccinellid larvae](#) have been observed feeding on aphid colonies.

Oleander aphids sequester cardiac glycosides, recognized heart poisons, from their host plants (Rothschild et al. 1970). They also fortify their cornicle secretions with these bitter, poisonous chemicals. Their bright aposematic (warning) coloration and possession of toxins protects them from predation by certain species of birds and spiders (Malcolm 1986). Spiders that have the cornicle secretion applied to their mouthparts immediately retreat and attempt to clean them. These cardiac glycosides appear not to harm the

parasitoids and generalist insect predators associated with oleander aphid.



Figure 5. Winged adult oleander aphid, *Aphis nerii* Boyer de Fonscolombe, with hole through which the parasitoid, *Lysiphlebus testaceipes* (Cresson), emerged.  
Credits: Lyle Buss, University of Florida



Figure 6. Colony of oleander aphid, *Aphis nerii* Boyer de Fonscolombe, that has suffered heavy parasitism by *Lysiphlebus testaceipes* (Cresson).  
Credits: J. Castner, University of Florida



Figure 7. Syrphid fly larva feeding on nymph of oleander aphid, *Aphis nerii* Boyer de Fonscolombe.  
Credits: J. Castner, UF/IFAS



Figure 8. Syrphid pupa on scarlet milkweed inflorescence infested with oleander aphid, *Aphis nerii* Boyer de Fonscolombe.  
Credits: Lyle Buss, University of Florida

Insecticidal soaps and oils are often effective against soft-bodied insects such as aphids. Other insecticides are not necessary in the home garden environment given the high level of natural biological control, ease of killing aphids with insecticidal soaps and oils, and incompatibility of use on butterfly nectar or larval host plants.

## Selected References

Blackman RL, Eastop VF. 2000. Aphids on the World's Crops: an Identification and Information Guide. Wiley, New York.

Essig EO. 1958. Insects and Mites of Western North America. MacMillan Publishers, New York.

Hall RW, Ehler LE. 1980. Population ecology of *Aphis nerii* on oleander. *Environmental Entomology* 9: 338-344.

Malcolm SB. 1986. Aposematism in a soft-bodied insect: a case for kin selection. *Behavioral Ecology and Sociobiology* 18: 387-393.

Rothschild M, von Euw J, Reichstein T. 1970. Cardiac glycosides in the oleander aphid, *Aphis nerii*. *Journal of Insect Physiology* 16: 1141-1145.

Shelton, A. *Lysiphlebus testaceipes*, (Hymenoptera: Aphidinae). *Biological Control: A Guide to Natural Enemies in North America*. (9 May 2017).